



Delirium Toolkit:

DELIRIUM GERI-GUIDE

PRE AND POST LEARNING ASSESSMENT WITH ANSWER GUIDE

Recommendations:

- Utilize this learning assessment tool before and after review of the Delirium Geri-Guide interactive flow chart (<https://www.geriguides.com/>) to document training, learner understanding and to identify additional learning needs.
- Utilize as one component of a comprehensive strategy for prevention, early identification, and management of delirium. A brief overview of how to navigate through the Geri-Guide and additional delirium tools and resources is located at <https://quality.allianthealth.org/topic/delirium/>

Name of Individual completing the Pre-Learning Assessment: _____

Date of Pre-Learning Assessment: _____ ID Number (if required): _____

Pre-Learning Assessment

1. Which of the following statements reflect appropriate use of antipsychotics to treat delirium? A. Antipsychotics are effective in preventing delirium B. Antipsychotics may resolve, shorten the duration, or lessen the intensity of delirium C. Antipsychotics may increase the risk of death D. Antipsychotics should only be reserved for patients who are risking injury to self, others, or having severe, distressing psychotic symptoms	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
2. What causes delirium? A. Any medical insult B. History of dementia C. Regular bowel movements	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
3. What are the types of delirium? A. Hyperactive delirium B. Hypoactive delirium C. Mixed delirium D. A and B E. All of the above	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
4. Which of the following is least likely to cause delirium in a patient/resident? A. Pneumonia B. Opioid use after surgery C. Diphenhydramine (Benadryl) D. Constipation	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
5. Which feature must be present to diagnose delirium? A. Inattention B. Confusion C. Agitation D. Altered sleep cycle	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>

Pre-Learning Assessment Score: _____

Post-Learning Assessment

<p>1. Which of the following is least likely to cause delirium in a patient/resident?</p> <p>A. Diphenhydramine (Benadryl) B. Constipation C. Opioid use after surgery D. Pneumonia</p>	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
<p>2. What causes delirium?</p> <p>A. History of dementia B. Regular bowel movements C. Any medical insult</p>	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
<p>3. Which feature must be present to diagnose delirium?</p> <p>A. Altered sleep cycle B. Inattention C. Confusion D. Agitation</p>	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
<p>4. What are the types of delirium?</p> <p>A. Hypoactive delirium B. Hyperactive delirium C. Mixed delirium D. A and C E. All of the above</p>	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
<p>5. Which of the following statements reflects appropriate use of antipsychotics to treat delirium?</p> <p>A. Antipsychotics may resolve, shorten the duration, or lessen the intensity of delirium B. Antipsychotics are effective in preventing delirium C. Antipsychotics should only be reserved for patients who are risking injury to self, others, or having severe, distressing psychotic symptoms D. Antipsychotics may increase the risk of death</p>	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>

Post-Learning Assessment Score: _____

Pre-Learning Assessment Answer Guide

1. Which of the following statements reflect appropriate use of antipsychotics to treat delirium? A. Antipsychotics are effective in preventing delirium B. Antipsychotics may resolve, shorten the duration, or lessen the intensity of delirium C. Antipsychotics may increase the risk of death D. Antipsychotics should only be reserved for patients who are risking injury to self, others, or having severe, distressing psychotic symptoms	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input checked="" type="checkbox"/>	E <input type="checkbox"/>
2. What causes delirium? A. Any medical insult B. History of dementia C. Regular bowel movements	A <input checked="" type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
3. What are the types of delirium? A. Hyperactive delirium B. Hypoactive delirium C. Mixed delirium D. A and B E. All of the above	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input checked="" type="checkbox"/>
4. Which of the following is least likely to cause delirium in a patient/resident? A. Pneumonia B. Opioid use after surgery C. Diphenhydramine (Benadryl) D. Constipation	A <input type="checkbox"/>	B <input checked="" type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
5. Which feature must be present to diagnose delirium? A. Inattention B. Confusion C. Agitation D. Altered sleep cycle	A <input checked="" type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>

Post-Learning Assessment Answer Guide

1. Which of the following is least likely to cause delirium in a patient/resident? A. Diphenhydramine (Benadryl) B. Constipation C. Opioid use after surgery D. Pneumonia	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input checked="" type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
2. What causes delirium? A. History of dementia B. Regular bowel movements C. Any medical insult	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input checked="" type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
3. Which feature must be present to diagnose delirium? A. Altered sleep cycle B. Inattention C. Confusion D. Agitation	A <input type="checkbox"/>	B <input checked="" type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>
4. What are the types of delirium? A. Hypoactive delirium B. Hyperactive delirium C. Mixed delirium D. A and C E. All of the above	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input type="checkbox"/>	D <input type="checkbox"/>	E <input checked="" type="checkbox"/>
5. Which of the following statements reflects appropriate use of antipsychotics to treat delirium? A. Antipsychotics may resolve, shorten the duration, or lessen the intensity of delirium B. Antipsychotics are effective in preventing delirium C. Antipsychotics should only be reserved for patients who are risking injury to self, others, or having severe, distressing psychotic symptoms D. Antipsychotics may increase the risk of death	A <input type="checkbox"/>	B <input type="checkbox"/>	C <input checked="" type="checkbox"/>	D <input type="checkbox"/>	E <input type="checkbox"/>