

DROPLET PRECAUTIONS



(In addition to Standard Precautions) (If you have questions, ask nursing staff)

Everyone Must:



Clean hands when entering and leaving room

Wear mask



Doctors and Staff Must:

Wear eye protection with respiratory symptoms and standard precautions if contact with secretions likely.













DROPLET PRECAUTIONS

If patient has diarrhea and/or C. difficile add Contact Enteric Precautions

Display sign outside the door. Remove sign after room is terminally cleaned.

Common Conditions (refer to Facility Policy):

- Influenza
- Meningitis
- Pertussis
- Respiratory viruses
- Mumps

Dishes/Utensils:

No special precautions. Kitchenware sanitized in dishwasher.

Equipment and Supplies:

- · Only essential equipment in room.
- Use dedicated or disposable equipment when available.
- Clean and disinfect reusable equipment including intravenous pumps, cell phone or pagers (if used in room), and other electronics, supplies, and other equipment prior to removing from patient's room.
- Ensure blood pressure cuff and stethoscope are cleaned and disinfected between patients.

Linen Management:

Bag linen in patient's room.

Personal Protective Equipment:

Standard and Tear-away Gown	Three-part Gown
Put ON in this order:	Put ON in this order:
1. <u>Wash or gel hands</u>	1. <u>Wash or gel hands</u>
2. Gown (if needed)	2. Gown (if needed)
3. Mask	3. Mask
4. Eye cover (if needed)	4. Eye cover (if needed)
5. Gloves (if needed)	5. Gloves (if needed)
Take OFF & dispose in this order:	Take OFF & dispose in this order:
1. Gloves (if used)	1. Gown and Gloves at the same time (grab gown
2. Eye cover (if used)	and pull off gloves in one movement)
3. Gown (if used)	2. Eye cover (if used)
4. Mask	3. Mask
5. <u>Wash or gel hands</u> (even if gloves used)	5. <u>Wash or gel hands</u> (even if gloves used)

Private Room:

If not available, please follow facility policy when cohorting patients.

Room Cleaning:

Follow facility policy for Droplet Precautions disinfection and curtain change requirements.

Transport:

Essential transport only, Have patient wear a surgical mask. Clean and disinfect transport vehicle. Alert receiving department regarding patient's isolation precaution status.

Discontinue precautions as per Facility Policy or Infection Prevention and Control Team instructions.





