



QUICK GUIDE TO STANDARDS OF CARE FOR CHRONIC DISEASE SCREENING

Recommendations for Chronic Kidney Disease (CKD) Screening

Patient Risk Factors	Screening Test	Clinical Indicators of CKD
<ul style="list-style-type: none"> ○ Diabetes ○ Hypertension ○ Age 60 years or older ○ Family history of CKD, diabetes, or hypertension ○ History of acute kidney injury ○ *U.S. ethnic minority status 	Single Void Specimen for albumin-to-creatinine ratio (ACR) to detect albuminuria Serum creatinine to estimate glomerular filtration rate (GFR)	Present for three months or more: <ul style="list-style-type: none"> ○ ACR >30 mg/g ○ GFR <60 mL/min/1.73 m² ○ Abnormalities of kidney structure or function

*Black/African American, Hispanic/Latino, Asian American, Pacific Islander, American Indian, Alaska Native, Native Hawaiian, or Other Pacific Islander

Annually assess urinary albumin levels using single void specimen for albumin-to-creatinine ratio (UACR) and evaluate the estimated glomerular filtration rate (eGFR) of:

- All patients with Type 2 diabetes annually, regardless of treatment.
- Patients with Type 1 diabetes with a disease duration of ≥5 years.

Bi-annual monitoring of UACR and eGFR is recommended to guide treatment for patients with diabetes and urinary albumin ≥300 mg/g creatinine and/or an eGFR 30–60 mL/min/1.73 m².

Current Procedural Terminology/Health Care Procedure Coding System Codes for Chronic Kidney Disease-Related Preventative Services	
Diabetes screening	82947, 82950, 82951
Cardiovascular disease screening	80061, 82465, 83718, 84478
Intensive behavioral therapy (IBT) for cardiovascular disease (CVD)	G0446
Intensive behavioral therapy (IBT) for obesity	G0447
Medical nutrition therapy (MNT) services	97802, 97803, 97804, G0270, G0271
Alcohol misuse screening & counseling	G0442, G0443
Counseling to prevent tobacco use	99406, 99407

References:

1. National Kidney Foundation (NKF) CKD Reference Card: <https://www.kidney.org/sites/default/files/NKF-CKD-Card.pdf>